




Phytophthora Dieback Occurrence Report for Kwinana Freeway Upgrade Assessment

Prepared for Westport Roads IPT

September 2025



Client	Westport Roads IPT
Report name	Phytophthora Dieback Occurrence Report for Kwinana Freeway Upgrade Assessment

Version History

Author	Reviewer	Version	Version Number	Submission Date
S. Hewitt	D. Delaporte	Draft	0.1	29/08/2025
S. Hewitt	D. Delaporte	Draft	0.2	22/09/2025
D. Delaporte	-	Final	1.0	23/09/2025

© Glevan Consulting 2025

26 Forrest Street, Pinjarra WA 6208

P: 1300 453 826

E: mail@glevan.com.au

W: www.glevan.com.au

This report has been prepared in accordance with the scope of work agreed between Westport Roads IPT and Glevan Consulting and contains results and recommendations specific to the agreement. Results and recommendations in this report should not be referenced for other projects without the written consent of Glevan Consulting.

Procedures and guidelines stipulated in various manuals, particularly Phytophthora Dieback Interpreters Manual for lands managed by the Department (DBCA), are applied as the base methodology used by Glevan Consulting in the delivery of the services and products required by this scope of work. These guidelines, along with overarching peer review and quality standards ensure that all results are presented to the highest standard.

Glevan Consulting has assessed areas based on existing evidence presented at the time of assessment. The Phytophthora pathogen may exist in the soil as incipient disease. Methods have been devised and utilised that compensate for this phenomenon; however, very new centres of infestation, that do not present any visible evidence, may remain undetected during the assessment.

Executive Summary

Glevan Consulting was commissioned by West Port Roads IPT to conduct a Phytophthora Dieback assessment of remnant bushland along Kwinana Freeway spanning from Farrington Road in the north to Safety Bay Road in the south. Portions of the Project Area had previously been assessed by Glevan Consulting, with the most recent surveys taking place in 2013 and 2014. The Project Area totalled 347.62 hectares (ha).

The assessment began on the 7th of July and was finalised on the 13th of August by Shannon Hewitt of Glevan Consulting. The assessment was carried out in accordance with the Phytophthora Dieback Interpreter's Manual for lands managed by the Department of Biodiversity, Conservation and Attractions (DBCA) (Department of Parks and Wildlife, 2015).

Of the intact native vegetation, 41.94 ha (12.06 %) were classed as Uninfested with adequate amounts of susceptible species present. Permanently Uninterpretable Protectable vegetation accounted for 27.78 ha (7.99 %) of the Project Area due to the overall lack of sufficient indicators, with a further 1.51 ha (0.43 %) ha classed as Permanently Uninterpretable Unprotectable. This classification was used due to the presence of an Infested area upslope of the Permanently Uninterpretable vegetation. 268.43 ha was classified as Excluded due to the complete lack of natural vegetation often found in paddocks, residential areas and private businesses where natural vegetation has become degraded or completely cleared. The remaining 5.47 ha (1.57 %) was classified as Infested.

Three soil and tissue samples were taken throughout the course of the assessment, one of which returned a positive result for *Phytophthora cinnamomi*.

The validity of the hygiene boundaries mapped for this project is twelve months from the completion of the assessment.

Table of Contents

1	Introduction	7
2	Background – Phytophthora Species and Dieback Disease	9
3	Materials and Methods.....	12
3.1	Project Area	12
3.2	Assessment Method	13
3.3	Collection of Evidence of Phytophthora Dieback	14
3.4	Demarcation	15
3.5	Soil and Tissue Samples	16
3.6	Determining Protectable Areas	18
4	Results	19
4.1	Desktop Assessment	19
4.2	Phytophthora Dieback Distribution	20
4.3	Disease Symptoms and Expression	21
4.4	Permanently Uninterpretable Areas	21
4.5	Temporarily Uninterpretable Areas	22
4.6	Excluded Areas	22
4.7	Allocation of Categories	22
4.8	Sample Summary	23
5	Discussion	24
6	Bibliography	26
7	Appendices.....	27
7.1	Phytophthora Dieback Occurrence Map	27
7.2	Mapping Metadata	27
7.3	Shapefile Spatial Data	28

List of Figures

Figure 1 – Kwinana Freeway Upgrade Project Area location.	8
Figure 2 – Phytophthora Dieback Occurrence Map 1.....	29
Figure 3 – Phytophthora Dieback Occurrence Map 2.....	30
Figure 4 – Phytophthora Dieback Occurrence Map 3.....	31
Figure 5 – Phytophthora Dieback Occurrence Map 4.....	32
Figure 6 – Phytophthora Dieback Occurrence Map 5.....	33
Figure 7 – Phytophthora Dieback Occurrence Map 6.....	34
Figure 8 – Phytophthora Dieback Occurrence Map 7.....	35

List of Tables

Table 1 – Keighery Vegetation Condition Scale and Assessability (Department of Parks and Wildlife, 2015).	12
Table 2 – Phytophthora Dieback Assessment for Vegetation Condition.....	13
Table 3 – Demarcation Flagging Colours.	16
Table 4 – Determination of Requirement for Sampling (Department of Parks and Wildlife, 2015).	17
Table 5 – Results Summary of the Project Area.	23
Table 6 – Soil and Tissue Sample Results.....	23

Glossary

Assessment Area	The portion of the Project Area where Phytophthora occurrence assessment is possible or will be possible after vegetation recovery.
Biomass	The total quantity or weight (density) of organisms in a given area.
Buffer	The area between the edge of visible disease symptoms and demarcation.
Clean On Entry	A requirement at a defined, signposted point where entering machinery and vehicles are to be free of soil, plant, and other material to minimise the risk of spreading weeds, pests and diseases.
DBCA	Department of Biodiversity, Conservation and Attractions.
Demarcation	The physical installation and representation of boundaries between hygiene categories after interpretation.
Phytophthora Dieback	The disease of plants caused by infection by the soil-borne organisms of the genus <i>Phytophthora</i> , of which <i>P. cinnamomi</i> is the most widespread and destructive.
Disease Impact	The degree of harm caused by Phytophthora Dieback (high, moderate, low).
Excluded Area	An area of high disturbance where natural vegetation is unlikely to recover.
Host	A plant species that may be Infested with the pathogen but not cause disease.
Indicator Species	Plant species susceptible to Phytophthora disease and reliably show early symptoms.
Infection	The invasion of the disease to an individual plant and not the population.
Infestation	The invasion of the disease into a population of plants.
Infested	An area that a registered interpreter has determined is expressing disease symptoms that indicate the presence of the pathogen <i>Phytophthora cinnamomi</i> .
Incipient Disease	A disease that is not visibly symptomatic.
Interpretation	Determining disease presence or absence in natural ecosystems using observable factors.
Not Yet Resolved	Phytophthora occurrence diagnosis cannot be made because of inconsistent evidence.
Pathogen	Phytophthora species.
Phytophthora Occurrence Assessment	The entire scope of work that relates to the delivery of a Phytophthora occurrence map and report.
Permanently Uninterpretable	A natural area with inadequate visible symptoms present to make a diagnosis.
Project Area (Study Area)	The proponent's area of interest, including the area where disturbance activities could occur, including access and egress to activity areas.
Protectable Area	Portions of the Project Area over which hygiene management rules for the plant pathogen <i>Phytophthora</i> , including clean on entry, will apply. These areas are generally free of disease.
Susceptible	Likely or liable to be harmed by Phytophthora pathogen.
Uninfested	An area that a registered interpreter has determined may be free of plant disease symptoms that indicate the presence of the pathogen <i>Phytophthora cinnamomi</i> .
Temporarily Uninterpretable	A naturally vegetated area that has had disturbance and is likely to recover from that disturbance in the short term.
Unprotectable	A disease-free area that is likely to become infested within a given time.
Vector	Any agent that acts as a carrier or transporter.

1 Introduction

Glevan Consulting was commissioned by Westport Roads IPT to conduct a Phytophthora Dieback assessment to determine the disease status of land associated with the Kwinana Freeway upgrade. The Project Area totalled 347.62 hectares (ha), spanning from Farrington Road in the north to Safety Bay Road in the south.

The Phytophthora Dieback occurrence assessment was completed in August 2025 by Miss Shannon Hewitt. Miss Hewitt is accredited by the Department of Environment and Conservation in the detection, diagnosis and mapping of the Dieback disease (Interpreter Registration No: DPW PDI 45). This accreditation recognises the skills and experience of Miss Hewitt. All Dieback detection, diagnosis and mapping were completed in accordance with *FEM047 Phytophthora Dieback Interpreter's Manual for Lands Managed by the Department* (Department of Parks and Wildlife, 2015).

The validity of the hygiene boundaries mapped for this project is twelve months from the completion of the assessment. The Phytophthora Dieback occurrence categories will expire August 2026.




Figure 1 – Kwinana Freeway Upgrade Project Area location.

2 Background – Phytophthora Species and Dieback Disease

Thousands of Australian native plant species are susceptible to Phytophthora Dieback (Dieback), a disease caused by the pathogen *Phytophthora cinnamomi* and other *Phytophthora* species. Whilst many root pathogens that cause disease are present within the Australian landscape, *Phytophthora cinnamomi* has caused the most significant impact to date and poses the biggest threat (Commonwealth of Australia, 2018). This disease is a significant threat to Australia's biodiversity, placing important plant species at risk of death, local extirpation or even extinction. Its impact on plant communities can also result in major declines in some insect, bird and animal species due to the loss of shelter, nesting sites and food sources. Phytophthora Dieback can cause permanent damage to ecosystems. Once an area is Infested with the pathogen, eradication is usually impossible. Awareness that human activity can easily spread the pathogen will help prevent an increase in the extent of this disease (Commonwealth of Australia, 2018).

P. cinnamomi is one of at least 32 *Phytophthora* species that exist in Australia. Whilst *P. cinnamomi* has to date had the largest impact on the Western Australia vulnerable zone, other destructive *Phytophthora* species are emerging in Australia such as *P. arenaria*, *P. constricta*, *P. cryptogea*, *P. elongata*, *P. gregata*, *P. megasperma* and *P. multivora*. *P. multivora* is notable in having been linked to declines in vegetation within the Western Australian vulnerable zone (Commonwealth of Australia, 2018).

P. cinnamomi is a microscopic water mould that belongs to the class Oomycetes. Oomycetes organisms are filamentous and absorptive and reproduce both sexually and asexually. *Phytophthora* species are considered parasitic, behaving largely as a necrotrophic pathogen causing damage to the host plant's root tissues because of infection and invasion (Department of Parks and Wildlife, 2015). The pathogen infects a host when it enters at a cellular level and damages the cell structure.

Dieback is the result of interaction between three physical components forming a 'disease triangle': the pathogen (*Phytophthora* species), the environment and the host. All three components are needed for the disease to develop over time. A population of hosts is made up of susceptible, infected and immune or resistant individuals.

The relationship between the presence of *Phytophthora* and the development of Dieback disease is variable based on the susceptibility of native plant species and the different environmental characteristics, landform types and rainfall zones across bioregions.

The plant host is a highly variable part of the disease triangle. Assessment sites may range from having no susceptible hosts present through to almost all plants present in the environment demonstrating high levels of susceptibility to *P. cinnamomi*. Over 2000 potential host species for the *P. cinnamomi* pathogen exist in Western Australia alone (Commonwealth of Australia, 2018). Skilled Dieback Interpreters are able to observe and interpret disease presence using hundreds of different susceptible plant species (referred to as indicator species) across multiple vegetation communities. Interpreters are able to recognise when disease detection is not possible due to insufficient hosts being present within an area.

Within the vulnerable zone of Western Australia, three main family groups are regarded as highly susceptible to Phytophthora Dieback disease:

1. Proteaceae
2. Ericaceae
3. Xanthorrhoeaceae.

The Proteaceae plant family demonstrates a high level of susceptibility to the *Phytophthora cinnamomi* pathogen. Genera such as *Banksia*, *Isopogon*, *Adenanthos*, *Persoonia*, *Petrophile* and *Xylomelum* demonstrate high susceptibility, providing reliable indicators through much of the vulnerable zone of Western Australia. *Andersonia*, *Astroloma* and *Leucopogon* species from the Ericaceae family also demonstrate high levels of susceptibility, whilst other species within the family do not demonstrate the same level of vulnerability to the pathogen. All species contained in the Xanthorrhoeaceae family are considered to be reliable indicator species.

Not all species within the vulnerable zone of Western Australia are susceptible to Dieback, with some species demonstrating a resistance to the pathogen. Resistance to the pathogen is recognised within genera such as *Acacia*, *Calothamnus*, *Eucalyptus* (excluding *Eucalyptus marginata*), *Lepidosperma* and *Melaleuca*, among others. Resistance to *P. cinnamomi* is dependent on a multitude of factors such as the ability for a species to compartmentalise the infected part of a root or the plants ability to quickly regrow new roots in order to replace those infected by the pathogen (Commonwealth of Australia, 2018). It has been recognised under field conditions that herbaceous perennials, annuals and

geophytes demonstrate greater resistance to Phytophthora Dieback compared to woody perennials (Commonwealth of Australia, 2018).

Where species considered to be resistant to the *P. cinnamomi* pathogen are observed to be dying suddenly, Armillaria Rot Disease (ARD) may be considered to be the cause of death. ARD is caused by the *Armillaria luteobubalina* fungus. Drought conditions are another contributing cause of death other than ARD to Phytophthora resistant species.

Temperature, rainfall and soil type all affect the impact that Dieback has on an area. *P. cinnamomi* does not generally cause significant vegetation declines in areas that receive less than 400 millimetres of rain annually and occur north of latitude 30° (O’Gara et al., 2005 as cited in Commonwealth of Australia, 2018). Within Western Australia, the vulnerable zone exists in areas of Mediterranean climate (warm to hot, dry summers and mild to cool, wet winters) where annual rainfall exceeds 400 millimetres in the southwest of Western Australia. The most significant impact of the pathogen is observed in areas receiving an annual rainfall of above 800 to 1000 millimetres.

Soil characteristics also play a significant role in the occurrence of Dieback. Moisture is essential for the survival of the *Phytophthora* pathogen and for sporangia production, with wet soil conditions and warm temperatures favouring the release of motile zoospores from the sporangia. Optimal temperatures for the survival and reproduction of the pathogen are between 21°C to 30°C, a temperature range that is common through the vulnerable zone of Western Australia (Department of Parks and Wildlife, 2015). The growth and reproduction of the pathogen is also affected by soil pH, with a favourable pH range of between five and six (mildly acidic). Strong host resistance is observed in areas with a high level of soil fertility, creating a less favourable soil environment for the pathogen.

Whilst autonomous (active) spread of the disease does occur, *P. cinnamomi* is passively spread (or vectored) by animals and human activity via the movement of soils, allowing the rapid infestation of a new area by the pathogen. The movement of even small pieces of vegetation or soil containing viable inoculum (zoospores, chlamydospores, oospores) can result in the spread of the pathogen. Passive spread by a vector can occur both overland and in subsurface water flow (Commonwealth of Australia, 2018). Human induced spread of the *P. cinnamomi* pathogen is a significant risk in the Western Australian landscape. Activities such as road construction and maintenance, commercial activities (timber harvesting and mining), off-road vehicle movement and bush walking activities can result in the spread of the disease should movement of Infested soil occur.

3 Materials and Methods

3.1 Project Area

Areas within the Project Area were categorised as Excluded if the vegetation was suffering from significant disturbance. Significant disturbance is based on Vegetation Condition Scale (Keighery, 1994) shown in Table 1. Any remaining area was categorised post-assessment into Phytophthora Dieback occurrence categories (Table 2).

Table 1 – Keighery Vegetation Condition Scale and Assessability (Department of Parks and Wildlife, 2015).

Assessability	Scale		Vegetation Condition
Assessable	1	Pristine	Pristine or nearly so; no obvious signs of disturbance.
	2	Excellent	Vegetation structure intact; disturbance affecting individual species and weeds are non-aggressive species.
	3	Very good	Vegetation structure altered, obvious signs of disturbance. For example, disturbance to vegetation structure caused by repeated fires, the presence of some more aggressive weeds, dieback, logging and grazing.
Possibly assessable – Discretion required	4	Good	Vegetation structure significantly altered by very obvious signs of multiple disturbances. Retains basic vegetation structure or ability to regenerate it. For example, disturbance to vegetation structure caused by very frequent fires, the presence of some very aggressive weeds at high density, partial clearing, dieback and grazing.
Not assessable – Excluded from assessment	5	Degraded	Basic vegetation structure severely impacted by disturbance. Scope for regeneration but not to a state approaching good condition without intensive management. For example, disturbance to vegetation structure caused by frequent fires, the presence of very aggressive weeds, partial clearing, dieback and grazing.
	6	Completely degraded	The structure of the vegetation is no longer intact, and the area is completely or almost completely without native species. These areas are often described as 'parkland cleared' with the flora comprising weed or crop species with isolated native trees or shrubs.

Table 2 – Phytophthora Dieback Assessment for Vegetation Condition.

Vegetation Condition	Phytophthora Occurrence Category
Naturally vegetated areas. Keighery disturbance rating of 3 or less. Phytophthora occurrence categorisation is possible.	Infested - Determined to have plant disease symptoms consistent with the presence of <i>Phytophthora cinnamomi</i> .
	Uninfested - Determined to be free of plant disease symptoms that indicate the presence of <i>P. cinnamomi</i>
	Permanently Uninterpretable - Undisturbed areas where susceptible plants are absent, or too few to make a determination of the presence or absence of <i>P. cinnamomi</i> .
	Not yet resolved.
Vegetation structure temporarily altered.	Temporarily Uninterpretable - Areas of disturbance where natural vegetation is likely to recover.
Vegetation structure severely altered. Keighery disturbance rating 4 or greater. Phytophthora occurrence assessment is not possible	Excluded.

3.2 Assessment Method

Prior to assessment, all information relevant to the assessment was assembled to assist the interpretation process. This information included previous assessments of the area, historic sample results, history of burning and possible other disturbances. This process is referred to as the Desktop Assessment.

All Phytophthora Dieback detection, diagnosis and mapping are performed to standards and procedures defined in *FEM047 Phytophthora Dieback Interpreter's Manual for lands managed by the department* (Department of Parks and Wildlife, 2015). These procedures are grounded on the presence of indicator species in the vegetation, and the observance of deaths in these plants. An indicator species is a plant species that is reliably susceptible to *Phytophthora cinnamomi*. Indicator Species Deaths (ISDs) alone do not necessarily indicate disease presence, and it is necessary to consider all environmental and ecological factors that may be present. These other factors include:

- Chronology of deaths,
- Pattern of deaths,
- Topographical position,
- Vectoring – causal agencies, and

- Biomass and biological diversity reduction.

Other causes of plant deaths need to be considered when determining the presence of Phytophthora Dieback (Department of Parks and Wildlife, 2015), including:

- Armillaria Root Disease,
- Various cankers,
- Insects,
- Drought, wind scorch and frost,
- Salinity and waterlogging,
- Fire and lightning,
- Senescence and competition,
- Physical damage, and
- Herbicides and chemical spills.

The Phytophthora Dieback assessment process is undertaken using the following basic sequence as documented in *FEM047 Phytophthora Dieback Interpreter's Manual for lands managed by the department* (Department of Parks and Wildlife, 2015):

- Observe,
- Hypothesise,
- Undertake evidence collection,
- Form a diagnosis,
- Test the diagnoses (sample results, previous assessment, second assessment),
- Evidence is reviewed with a new hypothesis formed, if necessary,
- Final field diagnoses determined.

The assessment was conducted using both comprehensive and linear assessment methodologies.

3.3 Collection of Evidence of Phytophthora Dieback

During the assessment process, the collection of evidence to support the field diagnosis was recorded using a tablet running the ESRI Field Maps application. Waypoints are recorded at locations to show evidence of:

- Where field diagnosis is certain or almost certain of Phytophthora Dieback infestation,
- Healthy indicator species where field diagnosis is almost certain of the site being Uninfested,

- Sites with too few or devoid of indicator species, thus supporting Permanently Uninterpretable classification, or
- Areas of disturbance, which are Temporarily Uninterpretable or Excluded.

Additional waypoints recorded include:

- Points requiring soil and tissue sampling,
- Points located where samples have been taken,
- Points located at indicator species deaths, and
- Points that need to be revisited for further examination.

Evidence recorded during a comprehensive assessment is collected along a series of transect lines spaced 50 meters apart, with evidence collected no more than every 100 meters along the transect lines. Linear assessments represented by an assessment area 25 meters either side of the centre line are assessed using a minimum of a single pass on each side of the centre line. Assessment areas that exceed 25 meters either side of the centre line are either assessed using the methodology of a comprehensive assessment or additional transect lines are established parallel to the centre line at increments of 50 meters.

Waypoint evidence, sample sites and results, and field demarcations are then digitised into a Phytophthora Dieback Occurrence Map according to the standards defined by the *FEM047 Phytophthora Dieback Interpreter's Manual for lands managed by the department* (Department of Parks and Wildlife, 2015).

3.4 Demarcation

The Phytophthora Dieback occurrence categories were demarcated in the field using the appropriate flagging tape (Table 3) tied to the vegetation at an appropriate buffer width. An appropriate buffer width is determined by Interpreters during the course of the assessment, and takes into consideration the following:

- Natural autonomous spread of the pathogen,
- Likely presence of incipient disease,
- Potential for surface water to flow into an area from an infestation,
- Current rate of spread,
- Location of natural drainage lines, such as creeks and brooks,

- Other site factors, including soil types that increase or decrease the rate of spread and degree of slope.

At a minimum, an up-slope buffer of 15 meters and a down-slope buffer width of 25 meters will be applied.

Table 3 – Demarcation Flagging Colours.

Boundary	Tape Colour and Width
Phytophthora Dieback Infested	Fluoro pink 25 mm
Permanently Uninterpretable, Temporarily Uninterpretable and Excluded (when required)	Pink and black striped ('tiger tape') 25 mm
Permanently Uninterpretable against Excluded and Temporarily Uninterpretable	Double band pink and black tiger tape
Armillaria infestation	Not usually demarcated, but if required, double flagging of pink and black tiger tape and fluoro pink 25mm.

3.5 Soil and Tissue Samples

Soil and tissue samples taken during the assessment were to standards and prescriptions defined in *FEM047 Phytophthora Dieback Interpreter's Manual for lands managed by the department* (Department of Parks and Wildlife, 2015). While species specific sampling techniques were undertaken, the general process to collect a soil and tissue sample is as follows:

- Tools and equipment are checked for the presence of soil and plant matter. Sterilisation will have occurred at the completion of the previous sample. Re-sterilise if required.
- Using the sampling tool (long or short handle mattock), dig down to expose the stem base and roots of the target plant, to a minimum depth of 30 cm.
- Remove sections of the root material and stem base from all sides of the target plant and place into a plastic sample bag.
- Collect several handfuls of soil from different depths around the base of the target plant and place into the plastic sample bag.
- Complete two metal tags with the sample tag information, including project name, sample number, company details, sample date and Interpreter initials. Place one tag into the bag.
- Complete a paper tag with the above information and secure to the sample bag.

- Mark the location of the sample with flagging tape and secure the second metal information tag to a plant within the immediate vicinity of the flag.
- Record the sample location and required information as an evidence point using ESRI Field Maps.
- Remove any material from the sampling tools and sterilise.
- Store the sample(s) in a cool location away from direct sunlight until transported to the laboratory.

All samples were analysed in the Vegetation Health Services (DBCA) laboratory using best-practice techniques.

Taking a soil and tissue sample from dead and dying plants is an integral part of assessment – although in some cases sampling is not essential. Sample results provide evidence to support field diagnostic decisions. The following table (Table 4) shows the need for sampling to assist the disease diagnosis process.

Table 4 – Determination of Requirement for Sampling (Department of Parks and Wildlife, 2015).

Observable factors indicating likelihood of <i>Phytophthora cinnamomi</i> presence				
ISD type	Multiple	Cluster	Scattered	Isolated
Species	Some or most indicator species	Any indicator plant	Any indicator plant	Any indicator plant
Pattern development	Obvious			Not obvious
Chronology	Obvious			Not obvious
Topographic situation	Gully/flat	Lower to mid slope	Mid slope to upper slope	Ridge
Causal agent	Obvious			Not obvious
Requirement for soil and tissue sample	Low	High	High	Low

Samples may also be taken for the following strategic reasons:

- Supporting Infested field diagnosis,
- Incipient, subtle or cryptic disease in apparent Uninfested sites, or
- Altering mapped Infested area boundaries.

3.6 Determining Protectable Areas

Following the determination of Phytophthora Occurrence categories, all interpretable vegetation was assessed for protectability, using the Department of Biodiversity, Conservation and Attractions (DBCA) Protectable Areas criteria.

The *FEM047 Phytophthora Dieback Interpreter's Manual for lands managed by the department* (Department of Parks and Wildlife, 2015) defines 'Protectable Areas' as those that:

- Have been determined to be free of the pathogen *Phytophthora* spp. by a registered Dieback Interpreter (all susceptible indicator plant species are healthy and no plant disease symptoms normally attributed to *Phytophthora* Dieback are evident),
- Consists of areas where human vectors are controllable (e.g., not an open road, private property), and
- Are positioned in the landscape and are of sufficient size (e.g., > 4 ha with axis >100 m) such that a qualified Interpreter judges that the pathogen will not autonomously engulf them in the short term (a period of a few decades), or
- Includes areas of high conservation and/or socio-economic value (for example, a small Uninfested area with a known population of a susceptible species of Threatened flora).

Areas deemed Unprotectable are displayed on the Phytophthora Dieback Occurrence Map (See Section 7.1) as grey hashed areas. By default, all areas not displayed as Unprotectable on the Phytophthora Dieback Occurrence Map have been categorised as Protectable.

4 Results

4.1 Desktop Assessment

Landforms and Vegetation

Vegetation mapping was conducted for the Project Area by GHD in 2025 (GHD, 2025). The report produced by GHD indicates that a number of reliable indicator species are expected to occur within the intact vegetation within the Project Area. It is expected that the Project Area will offer good Phytophthora Dieback disease expression if *P. cinnamomi* were present.

Fire History

Analysis of the DBCA Fire History (DBCA-060) database (Department of Biodiversity, Conservation and Attractions, 2025) indicates that multiple fires have been recorded along the Project Area, two of these being wildfires that occurred in 2015 and 2021. These fires were located approximately one kilometre south of Anketell Road. Sufficient indicator species regeneration is expected to have occurred since the fires to give an accurate Dieback assessment.

Historical Assessments

Review of Glevan Consulting's database and publicly available data from the Phytophthora Dieback Occurrence - Infested Only (DBCA-082) dataset (Department of Biodiversity, Conservation and Attractions, 2025) does show previous surveys along and around the current Kwinana Freeway Project Area.

The closest positive samples can be found:

- A single positive soil and tissue sample was located within the project boundary on the eastern side of the Armadale Road southbound ramp.
- 420 metres southeast of the most eastern side of the Project area on Roe Highway, this positive can be found on the road Training Place.
- 240 metres west of the Project Area, found in between Berrigan Drive and Verna Court.

Additionally, the City of Kwinana advised representatives from GHD that Phytophthora Dieback is known to occur within a Lot south of Karakin Place.

Rainfall

The long-term annual average rainfall (2002 to 2025) recorded at the nearby Anketell Weather Station is 771.8 mm (Bureau of Meteorology, 2025), which means the Project Area is located in the 600 mm and above zone, where Phytophthora Dieback occurrence would not be restricted and would be expected to occur across the landscape.

4.2 *Phytophthora Dieback Distribution*

Two areas of Phytophthora Dieback disease were observed within the Project Area. The first area is located approximately 250 metres south of Gibbs Road on the eastern side of the Kwinana Freeway (Figure 4). A positive soil and tissue sample result was obtained in this area. The sample was located at the top of a hill, with downhill slopes on the north, south and west sides.

The second infestation is located north of North Lake Road to the west of Kwinana Freeway (Figure 3). Soil and tissue sampling was not possible in this area due to buried gas services. While a sample could not be taken during this assessment, a historic positive soil and tissue sample was observed to occur on the opposite side of Kwinana Freeway. The presence of visible symptoms of disease and the historic positive provided sufficient evidence to classify the area as Infested.

There were no observable symptoms of Phytophthora Dieback within the Project boundary adjacent to Karakin Place due to the vegetation being classified as Permanently Uninterpretable due to a lack of Indicator Species.

Vegetation classed as Uninfested (41.94 ha; 12.06 %) appeared to be in good condition with scattered indicators deaths seen throughout. These deaths were often found on slopes and ridges where there is a higher risk of drought and were thus classed as background deaths. Indicators throughout the assessment include:

- *Banksia attenuata*,
- *Banksia menziesii*,
- *Daviesia* species,
- *Lomandra* species,
- *Patersonia* species,
- *Stirlingia latifolia*,
- *Styphelia* species, and

- *Xanthorrhoea preissii*.

The remaining bushland and cleared bushland within the assessment categorised as Permanently Uninterpretable, Unprotectable Uninterpretable, Temporarily Uninterpretable and Excluded.

4.3 Disease Symptoms and Expression

Disease expression varied in the area where the positive soil and tissue sample result was recovered. A wetland was located towards the southern end of the infestation, with deaths of *Banksia attenuata* and *Xanthorrhoea preissii* observed, displaying perceptible expression. Scattered *Banksia attenuata* deaths were observed throughout the Sheoak (*Allocasuarina fraseriana*) without any discernible chronology and represented subtle disease expression. Old sparse *Xanthorrhoea preissii* deaths were also found across the slope.

4.4 Permanently Uninterpretable Areas

Several areas were classified as Permanently Uninterpretable due to insufficient coverage of reliable indicator species. Vegetation mapping conducted by GHD (2025) determined these areas to be *Corymbia calophylla* woodland (Cc), *Eucalyptus rudis* and *Melaleuca raphiophylla* woodland (ErMr) and roadside revegetation mapped as 'Isolated *Corymbia calophylla*', 'Mixed Shrubland', or 'Isolated *Eucalyptus* spp.'. These areas contained only sparse occurrences of reliable indicator species, limiting the ability to accurately assess disease presence or absence.

Permanently Uninterpretable vegetation accounted for 29.29 ha (8.42 %) of the total Project Area. Of this, 27.78 ha was considered Protectable from Phytophthora Dieback. These areas were considered Protectable due to occurring upslope from known infestations or being of sufficient size to not be consumed by Phytophthora Dieback in the short term. The remaining 1.51 ha of Permanently Uninterpretable vegetation occurred downslope of a new infestation identified during the current survey and will likely be consumed within the short term due to autonomous movement of the pathogen.

4.5 Temporarily Uninterpretable Areas

One section of the Project Area approximately 160 metres was classified as Temporarily Uninterpretable on the western side of Kwinana Freeway. This length can be found just south of Gaebler Road and approximately 760 metres north of Rowley Road. No current data was available for this portion on the DBCA Fire History (DBCA-060) database (Department of Biodiversity, Conservation and Attractions, 2025). After an infield assessment of the area, the fire was estimated to have occurred one to two years prior, resulting in insufficient indicators to accurately determine the disease status. It is usually estimated that a fire affected area will return to an interpretable status within three to five years after the initial burn, with wildfires and prescribed burns potentially affecting the recovery time based on the temperature of the burn.

4.6 Excluded Areas

Excluded areas are not able to be assessed for the presence of Phytophthora Dieback due to the level of disturbance of the vegetation, based on the Keighery Vegetation Condition Scale. A total of 268.43 ha of land within the Project Area has been classed as Excluded during the assessment. Of this, 4.36 ha was determined to be Unprotectable due to adjoining Phytophthora Dieback infestations.

4.7 Allocation of Categories

The Phytophthora Dieback Occurrence categorised mapped during the assessment are tabulated in Table 5.

Table 5 – Results Summary of the Project Area.

Category	Area (ha)	% of total area
Infested	5.47	1.57
Uninfested Protectable	41.94	12.06
Permanently Uninterpretable Protectable	27.78	7.99
Permanently Uninterpretable Unprotectable	1.51	0.43
Temporarily Uninterpretable Protectable	2.50	0.72
Excluded Protectable	264.07	75.96
Excluded Unprotectable	4.36	1.25
TOTAL AREA	347.62	

4.8 Sample Summary

Three soil and tissue samples were taken during the course of the assessment (Table 6).

Table 6 – Soil and Tissue Sample Results.

Sample Number	Plant Sampled	Easting	Northing	Result
SH01	<i>Lomandra species</i>	391363	6448612	Negative
SH02	<i>Xanthorrhoea preissii</i>	392278	6441299	<i>P. cinnamomi</i>
SH03	<i>Banksia attenuata</i>	391616	6433957	Negative

5 Discussion

Two areas of *Phytophthora* Dieback infestation were identified within the Project Area. The first is located approximately 250 metres south of Gibbs Road on the eastern side of the Kwinana Freeway (Figure 4), where a positive result was obtained through soil and tissue sampling. This sample was taken from the top of a hill, with surrounding slopes extending north, south and west. Disease expression in this area was variable, with observable deaths of *Banksia attenuata* and *Xanthorrhoea preissii* near a large wetland to the south. Scattered deaths of *Banksia attenuata* within surrounding *Allocasuarina fraseriana* woodland were also noted, although these lacked a clear chronology and displayed subtle expression. Sparse and aged *Xanthorrhoea preissii* deaths were recorded across the slope.

The second infestation was identified north of North Lake Road on the western side of the Kwinana Freeway (Figure 3). Soil and tissue sampling could not be conducted in this location due to the presence of underground gas infrastructure. However, visible disease symptoms, combined with a historic positive sample located on the opposite side of the Freeway, provided sufficient evidence to classify this area as Infested.

Vegetation assessed as Uninfested appeared to be in generally good condition. Scattered deaths of indicator species were observed, particularly on elevated terrain such as ridges and slopes where drought stress is more likely. These were interpreted as background deaths unrelated to disease activity. Indicator species recorded throughout the assessment included *Banksia attenuata*, *Banksia menziesii*, *Daviesia* spp., *Lomandra* spp., *Patersonia* spp., *Stirlingia latifolia*, *Styphelia* spp., and *Xanthorrhoea preissii*.

A number of areas were classified as Permanently Uninterpretable due to the absence of sufficient coverage by reliable indicator species. Vegetation mapping conducted by GHD (2025) identified these areas as *Corymbia calophylla* woodland (Cc), *Eucalyptus rudis* and *Melaleuca raphiophylla* woodland (ErMr), and roadside revegetation units classified as 'Isolated *Corymbia calophylla*', 'Mixed Shrubland', or 'Isolated *Eucalyptus* spp.'. The dominance of native wetland species and the degraded nature of some roadside environments further reduced diagnostic confidence, as many of these species are either non-susceptible to *Phytophthora cinnamomi* or exhibit limited and unreliable expression of disease symptoms.

One section of the Project Area, approximately 160 metres in length, located south of Gaebler Road and 760 metres north of Rowley Road on the western side of the Kwinana Freeway, was classified as Temporarily Uninterpretable. No recent fire data was available for this location in the DBCA Fire History database, though in-field observations indicated a fire had occurred one to two years prior. Post-fire vegetation recovery was insufficient at the time of assessment, with limited indicator regrowth observed. Typically, such areas are expected to return to an interpretable condition within three to five years, although recovery rates vary depending on fire intensity and type.

A total of 268.43 ha was Excluded from the assessment due to the level of disturbance. These Excluded areas were either highly degraded, completely cleared, or contained insufficient remnant vegetation. Of this, 4.36 ha were considered Unprotectable due to their proximity to existing Dieback infestations, which reduces the feasibility of effective hygiene management.

The validity of the hygiene boundaries mapped for this project is twelve months from the completion of the assessment. The Phytophthora Dieback occurrence categories will expire August 2026.

6 Bibliography

-
- Bureau of Meteorology. (2025). *Monthly rainfall - Anketell*. Retrieved from Bureau of Meteorology: http://www.bom.gov.au/jsp/ncc/cdio/weatherData/av?p_nccObsCode=139&p_display_type=dataFile&p_startYear=&p_c=&p_stn_num=009258
- Commonwealth of Australia. (2018). *Threat abatement plan for disease in natural ecosystems caused by Phytophthora cinnamomi*.
- Department of Biodiversity, Conservation and Attractions. (2025). *DBCA Fire History (DBCA-060)*. Retrieved from DataWA: <https://catalogue.data.wa.gov.au/dataset/dbca-fire-history>
- Department of Biodiversity, Conservation and Attractions. (2025). *Phytophthora Dieback Occurrence - Infested Only (DBCA-082)*. Retrieved from DataWA: <https://catalogue.data.wa.gov.au/dataset/phytophthora-dieback-occurrence-infested-only-dbca>
- Department of Parks and Wildlife. (2015). *FEM047 Phytophthora Dieback Interpreter's Manual for lands managed by the department*. Unpublished.
- GHD. (2025). *Kwinana Freeway Widening Project*. Unpublished report for Main Roads Western Australia.
- Keighery, B. (1994). *Bushland Plant Survey: a Guide to Plant Community Survey for the Community*. Wildflower Society of WA (Inc.).

7 Appendices

7.1 *Phytophthora Dieback Occurrence Map*

The provided map is the *Phytophthora Dieback* occurrence map.

The assessment area is displayed as a blue boundary line. The following categories are also shown:

- Excluded (shown as yellow). Areas of high disturbance where natural vegetation has been cleared and is unlikely to recover to a level that is interpretable.
- Infested (shown as red). Determined from the assessment to have the plant disease caused by *Phytophthora cinnamomi*. *Phytophthora Dieback* caused by other *Phytophthora* species will be displayed as other colours, typically shades of orange and yellow.
- Uninfested (shown as green). Determined from the assessment to be free of plant disease *Phytophthora Dieback*.
- Permanently Uninterpretable (shown as purple). Undisturbed areas where susceptible plants are absent, or too few to decide the presence or absence of *Phytophthora Dieback*.
- Temporarily Uninterpretable (shown as grey). Areas of disturbance where natural vegetation is likely to recover.

Additional spatial data that may be shown include:

- Sample location.

7.2 *Mapping Metadata*

Dataset Description	
Title	Kwinana_Freeway_Upgrades_2025
Data Created	29/08/2025
Date Last Updated	29/08/2025
Abstract	Phytophthora Dieback occurrence, samples, evidence, and project area shapefiles for the Kwinana Freeway Upgrade assessment.
Purpose	Dieback category boundary mapping.
Document Number	GC-25-0030
Contact Organisation	Glevan Consulting
Contact Name	Shannon Hewitt
Contact Position	Dieback Interpreter
Contact Phone	0487 253 158
Contact Email	shannon.hewitt@glevan.com.au

Lineage	All field data recorded using ESRI Field Maps application on a GPS enabled tablet.
Datum / Coordinate System	GDA2020 MGA Zone 50
Geographic Description	Kwinana, Western Australia
Restrictions	None

7.3 Shapefile Spatial Data

Shapefile Spatial Data	
File Contents	File Name
Occurrence	Kwinana_Freeway_Upgrades_2025_Occurrence
Samples	Kwinana_Freeway_Upgrades_2025_Samples
Evidence	Kwinana_Freeway_Upgrades_2025_Evidence
Project Area	Kwinana_Freeway_Upgrades_2025_Project_Area




Figure 2 – Phytophthora Dieback Occurrence Map 1.




Figure 3 – Phytophthora Dieback Occurrence Map 2.





Figure 4 – Phytophthora Dieback Occurrence Map 3.






Figure 6 – Phytophthora Dieback Occurrence Map 5.




Figure 7 – Phytophthora Dieback Occurrence Map 6.




Figure 8 – Phytophthora Dieback Occurrence Map 7.



GLEVAN
CONSULTING